

A REGIONAL COMPARISON OF CHILDREN'S BLOOD CADMIUM, LEAD, AND MERCURY IN RURAL, URBAN AND INDUSTRIAL AREAS OF SIX EUROPEAN COUNTRIES, AND CHINA, ECUADOR, AND MOROCCO

FRANTIŠKA HRUBÁ¹, MILENA ČERNÁ², CHUNYING CHEN³, FLORENCIA HARARI⁴, MILENA HORVAT⁵, KVETOSLAVA KOPPOVÁ⁶, ANDREA KRŠKOVÁ⁷, JAWHAR LAAMECH⁸, YU-FENG LI⁹, LINA LÖFMARK¹⁰, THOMAS LUNDH¹⁰, BADIYA LYOUSSI¹¹, DARJA MAZEJ⁵, JOŠKO OSREDKAR¹², KRYSZYNA PAWLAS¹³, NATALIA PAWLAS¹⁴, ADAM PROKOPOWICZ¹⁵, GERDA RENTSCHLER¹⁰, JANJA SNOJ TRATNIK⁵, JOHAN SOMMAR¹⁶, VĚRA SPĚVÁČKOVÁ⁷, ZDRAVKO ŠPIRIĆ¹⁷, STAFFAN SKERFVING¹⁰, and INGVAR A. BERGDAHL¹⁶

¹ Regional Authority of Public Health, Banská Bystrica, Slovakia

² Charles University, Prague, Czech Republic

Third Faculty of Medicine

³ Chinese Academy of Sciences, Beijing, China

Institute of High Energy Physics and National Center for Nanoscience and Technology

⁴ Institute for Development of Production and Work Environment (IFA), Quito, Ecuador

⁵ Institut Jožef Stefan, Ljubljana, Slovenia

Department of Environmental Sciences

⁶ Slovak Medical University Bratislava, Banská Bystrica, Slovakia

Faculty of Health

⁷ National Institute of Public Health, Prague, Czech Republic

Funding: this study was supported by the European Union's Sixth Framework Programme (grant No. FOOD-CT-2006-016253 entitled "PHIME – Public Health Impact of Long-Term, Low-Level Mixed Element Exposure in Susceptible Population Strata," coordinator: Staffan Skerfving), the Swedish Environment Protection Agency (grant No. dnr 721-1510-07 entitled: "Time series – lead concentrations in blood," grant manager: Staffan Skerfving, and Dnr 235-1451-10Mm entitled, "Lead, cadmium and mercury in blood of Skåne children 2009-2011 – with comparisons back to 1978," grant manager: Staffan Skerfving), the Polish Ministry of Science and Higher Education (decision No. 193/6. PR UE/2007/7 entitled "Public health impact of long-term, low-level mixed element exposure in susceptible population strata," project manager: Krystyna Pawlas), the National Institute of Public Health in Prague and the Ministry of Health of the Czech Republic (grant No. 016253 entitled "Public health impact of long-term, low-level mixed element exposure in susceptible population strata," project manager: prof. MUDr. Milena Černá, DrSc.), Public Health Authority of the Slovak Republic (project of the Authorities of Public health SR : 1.9 entitled „Public health impact of long-term, low-level mixed element exposure in susceptible population strata," project manager: Kvetoslava Koppova), and the Slovenian Research Agency (ARRS) (program No P1-0143 entitled „Cycling of substances in the environment, mass balances, modelling of environmental processes and risk assessment," program manager: Milena Horvat).

The paper reflects only the authors' views; the European Union is not liable for any use that may be made of the information.

Received: January 7, 2023. Accepted: May 30, 2023.

Corresponding author: Ingvar A. Bergdahl, Umeå University, Department of Public Health and Clinical Medicine, Division of Sustainable Health, 90187 Umeå, Sweden (e-mail: ingvar.bergdahl@umu.se).

⁸ Abdelmalek Essaadi University, Tangier, Morocco

Laboratory of Chemistry and Biochemistry, Faculty of Medicine and Pharmacy

⁹ Chinese Academy of Sciences, Beijing, China

Institute of High Energy Physics

¹⁰ Lund University, Lund, Sweden

Division of Occupational and Environmental Medicine

¹¹ University Sidi Mohamed Ben Abdellah, Fez, Morocco

Laboratory of Natural Substances, Pharmacology, Environment, Modelling, Health and Quality of Life

¹² University Medical Centre Ljubljana, Ljubljana, Slovenia

¹³ Wrocław Medical University, Wrocław, Poland

Department of Hygiene

¹⁴ Medical University of Silesia, Zabrze, Poland

Department of Pharmacology, Faculty of Medical Sciences in Zabrze

¹⁵ Institute for Ecology of Industrial Areas, Katowice, Poland

¹⁶ Umeå University, Umeå, Sweden

Department of Public Health and Clinical Medicine, Division of Sustainable Health

¹⁷ Green Infrastructure Ltd, Zagreb, Croatia

Abstract

Objectives: The authors aimed to evaluate whether blood cadmium (B-Cd), lead (B-Pb) and mercury (B-Hg) in children differ regionally in 9 countries, and to identify factors correlating with exposure. **Material and Methods:** The authors performed a cross-sectional study of children aged 7–14 years, living in 2007–2008 in urban, rural, or potentially polluted (“hot spot”) areas (ca. 50 children from each area, in total 1363 children) in 6 European and 3 non-European countries. The authors analyzed Cd, Pb, and total Hg in blood and collected information on potential determinants of exposure through questionnaires. Regional differences in exposure levels were assessed within each country. **Results:** Children living near industrial “hot-spots” had B-Cd 1.6 (95% CI: 1.4–1.9) times higher in the Czech Republic and 2.1 (95% CI: 1.6–2.8) times higher in Poland, as compared to urban children in the same countries (geometric means [GM]: 0.13 µg/l and 0.15 µg/l, respectively). Correspondingly, B-Pb in the “hot spot” areas was 1.8 (95% CI: 1.6–2.1) times higher than in urban areas in Slovakia and 2.3 (95% CI: 1.9–2.7) times higher in Poland (urban GM: 19.4 µg/l and 16.3 µg/l, respectively). In China and Morocco, rural children had significantly lower B-Pb than urban ones (urban GM: 64 µg/l and 71 µg/l, respectively), suggesting urban exposure from leaded petrol, water pipes and/or coal-burning. Hg “hot spot” areas in China had B-Hg 3.1 (95% CI: 2.7–3.5) times higher, and Ecuador 1.5 (95% CI: 1.2–1.9) times higher, as compared to urban areas (urban GM: 2.45 µg/l and 3.23 µg/l, respectively). Besides industrial exposure, traffic correlated with B-Cd; male sex, environmental tobacco smoke, and offal consumption with B-Pb; and fish consumption and amalgam fillings with B-Hg. However, these correlations could only marginally explain regional differences. **Conclusions:** These mainly European results indicate that some children experience about doubled exposures to toxic elements just because of where they live. These exposures are unsafe, identifiable, and preventable and therefore call for preventive actions. *Int J Occup Med Environ Health.* 2023;36(3):349–64

Key words:

biological monitoring, child, mercury, lead, environmental pollutants, cadmium

INTRODUCTION

Cadmium, lead, and mercury are classic toxic elements. Exposure occurs through ingestion and/or inhalation, which, in turn, may depend on where we live [1–5]. For example, urban and rural areas may differ, and local contamination, industrial or other, can potentially affect exposure.

Lead affects the brain, especially its development. Associations between lead and intelligence and/or school performance have been found even at very low exposure levels, reflected by blood-lead concentrations [2,3,5]. For cadmium, recent studies suggest that the brain, the kidney and skeleton is affected [6,7]. No safe level

of exposure can be defined for lead, as there is no evidence for a threshold [2] and this may be the case also for cadmium [7], even when considering only non-cancer effects. There is, thus, a clear concern for health effects at the exposures that occur in the general population. For mercury, methylmercury from fish and elemental mercury from dental amalgam are health concerns [8]. There have been concerns that the expansion of heavy metallurgic industry in Eastern Europe during the 1950s and 1960s led to significant, still prevailing, exposure in, e.g., Poland and former Czechoslovakia.

In order to learn more about geographic differences of exposure to these elements, international harmonized human biomonitoring of children's cadmium, lead, and mercury concentrations in blood (B-Cd, B-Pb, and B-Hg, respectively) was made in 2007–2008 in 9 countries within the project Public Health Impact Of Long-Term, Low-Level Mixed Element Exposure In Susceptible Population Strata (PHIME). Six of the countries were European, mainly in Eastern Europe, the other 3 were in China, Ecuador, and Morocco. The authors have previously published international comparisons of results from blood samples from the urban children [9] and from women [10,11], analyzed in 1 single laboratory. These studies showed that the international differences between the studied European countries are modest for cadmium and lead, meaning that the urban children in the studied Eastern European countries did not have much higher exposures than children in Sweden. A later study of cadmium within the European human biomonitoring initiative HBM4EU similarly found no distinct geographic patterns for cadmium in adults' urine in an international comparison of 9 European countries [12], whereas the authors are unaware of any corresponding publications on lead or mercury. While thus European international differences in the general urban population appear to be modest, children may still be additionally exposed from local sources, and it is therefore important

not to neglect the issue of regional exposures. Urban and rural areas may differ and it is well-known that living close to an industry or workshop that handles lead, or an industrially contaminated area, can be a major source of lead exposure [5,13]. The authors therefore here further analyze regional variation within countries, including rural children and children from polluted areas, so called "hot spots," in order to learn more about how regional variation within countries affect children's B-Cd, B-Pb, and B-Hg. Concentrations were not compared between countries, only within countries, by reasons described in Material and Methods section.

Thus, the aim of the present study was to describe regional variation in children's B-Cd, B-Pb, and B-Hg between 3 areas in each of the 9 countries where urban children's levels have been previously compared. Also for the European countries, based on questionnaire data, some correlates with exposure, other than geography were described.

MATERIAL AND METHODS

Subjects and study areas

The study was performed in 9 countries: Croatia, the Czech Republic, Poland, Slovakia, Slovenia, Sweden, China, Ecuador, and Morocco, in 2007–2008 (Table 1). In each country, schools were selected in 3 different areas: urban, rural and industrial (potentially polluted "hot spots"). The industrial areas were selected on basis of local environmental researchers' knowledge or suspicion of former or current contamination with heavy metals. Contact was established with the management of each school, and about 50 school children aged 7–11 years (in Morocco up to 14) were recruited. Blood samples and questionnaire data were collected. Data from Morocco have been published separately in a study of renal biomarkers [14] and the urban children are the same as in the previous study [9]. The study was approved by the local ethical committees. Written consent was obtained from a parent of each child and oral consent from the child before sampling.

Table 1. Characteristics of the studied areas and the 1363 children examined 2007–2008, living in urban, rural, or potentially polluted areas in 6 European and 3 non-European countries

Area of sampling	Type of contamination	Participants (girls/boys) [n]	Age [years] (M (range))
Croatia			
urban (Koprivnica)		52 (27/25)	8.7 (8–10)
rural		56 (21/35)	9.0 (7–10)
industrial	petrochemical	42 (21/21)	8.7 (7–10)
Czech Republic			
urban (Prague)		25 (8/17)	8.3 (7–10)
rural		66 (31/35)	9.7 (8–11)
industrial	predominantly Pb/Cd smelter	97 (50/47)	8.9 (7–11)
Poland			
urban (Wroclaw)		35 (10/25)	8.0 (7–8)
rural		39 (22/17)	8.4 (8–9)
industrial	Pb smelter	43 (16/27)	8.5 (8–10)
Slovakia			
urban (Banská Bystrica)		57 (35/22)	8.9 (7–11)
rural		50 (18/32)	8.3 (7–10)
industrial	Hg smelter	50 (31/19)	8.6 (7–10)
Slovenia			
urban (Ljubljana)		45 (26/19)	9.0 (7–11)
rural		65 (31/34)	9.5 (6–11)
industrial	Hg mining	65 (30/35)	9.2 (8–11)
Sweden			
urban (Landskrona)		41 (19/22)	9.1 (8–11)
rural		48 (22/26)	9.1 (8–11)
industrial	Pb smelter	59 (31/28)	9.2 (8–11)
China			
urban (Guiyang)		50 (30/20)	8.2 (7–10)
rural		58 (33/25)	7.3 (7–8)
industrial	Hg mining	42 (15/27)	8.5 (7–10)
Ecuador			
urban (Camilo Ponce Enriquez)		68 (36/32)	7.2 (7–10)
rural		47 (28/19)	7.6 (7–9)
industrial	gold mining	32 (10/22)	7.8 (7–10)
Morocco			
urban (Fez, Sefrou)		39 (18/21)	10.1 (7–14)
rural		68 (38/30)	9.9 (7–14)
industrial	heavy industry (ceramics, metals)	24 (18/6)	8.9 (8–11)

The total number of children examined was 1363 (Table 1). Of the industrial areas/potential “hot spots,” 4 were situated around Pb/Cd smelters, 2 were Hg-mining locations, 1 – a gold mining community using Hg for refining of gold, 1 was a petrochemical area and 1 was a heavy industry area (both ceramics and metals). In Morocco, Pb could possibly still be used in petrol, at least to some extent. Therefore, Pb contamination was expected in Morocco’s urban areas.

Questionnaire/interview and sampling

Information on individual factors of potential concern for metals exposure was obtained through 3 sources:

- questionnaire to the school,
- questionnaire to the parents,
- interview with/examination of the child.

The questionnaires and interview forms were jointly developed for all countries and were translated into the local languages. Questions were identical, but local additions were allowed as long as they could not be expected to bias answers to the core questions. From the parents’ questionnaire, information on their education (basic, middle, high), whether anyone smoked at home (yes/no), source of water and heating, traffic near the home (<1 vs. ≥ 1 car passing/min), and the child’s intake of offal (<1 vs. ≥ 1 meal/month), fish (<1, 1–2, ≥ 3 meals/month), and shellfish (<1 vs. ≥ 1 meal/month) was obtained. Information on number of dental amalgam fillings and the child’s attempts to smoke was obtained at the examination.

A nurse took a cubital venous blood sample after cleaning with an ethanol swab. Evacuated plastic tubes with heparin (Greiner Vacuette® 4 ml Lithium Heparin tubes, Greiner-Bio One GmbH, Frickenhausen, Germany) were used. Sampling tubes from one and the same production batch were used in all participating countries. The levels of all 3 metals were $<0.03 \mu\text{g/l}$ at leaching tests of tubes with 4 ml of 2% nitric acid.

Chemical analyses

Analyses of B-Pb, B-Cd and B-Hg were carried out in 5 laboratories (the Czech Republic, Poland, Slovenia, Sweden, and China, which analyzed their own samples; Sweden also analyzed samples from Slovakia, Ecuador and Morocco; Slovenia also from Croatia). In addition, 1 laboratory (Sweden; central laboratory) re-analyzed B-Pb, B-Cd and B-Hg in the urban samples from all countries except those already analyzed in Sweden and those from China where only B-Hg was re-analyzed (because of limited remaining sample volumes), and Slovenia and Croatia where only B-Pb and B-Cd were re-analyzed [9]. Thus, many of the urban samples were analyzed twice. By choosing the Swedish laboratory as central laboratory the number of re-analyses was minimized, as samples from 4 countries had already been analyzed in that laboratory.

Blood cadmium and B-Pb were determined by inductively coupled plasma mass spectrometry (ICP-MS) (samples from Slovenia, Sweden, and China according to the methods modified from Barany et al. [15]) or electrothermal atomic absorption spectrometry (ETAAS, Poland; [16]; and the Czech Republic), B-Hg by ICP-MS (China), cold vapour AAS (Poland; [16]), cold vapour atomic fluorescence spectrometry (Sweden; [17]), or by thermal combustion, amalgamation and atomic absorption spectrometry using Direct Mercury Analyzer (DMA-80) (Slovenia; EPA method 7473), or the single-purpose Atomic Mercury Analyzer AMA 254 (the Czech Republic).

Although each laboratory used its internal quality assurance system (Table 2), involving the use of reference materials, and demonstrated good performance in external proficiency testing schemes, it was decided to organize 4 interlaboratory comparisons. The aim was to demonstrate the level of equivalence between the participating laboratories and to decide whether samples should be analyzed in separate laboratories or centrally. Three intercomparisons used lyophilised samples of human blood from:

Table 2. Quality control procedures reported from the laboratories participating in the study on children aged 7–14 years, living in 2007–2008 in urban, rural, or potentially polluted areas in 6 European and 3 non-European countries


Country	Reference material	Intercomparisons	Accreditation
Sweden	Seronorm whole blood level 1	Centre de Toxicologie du Quebec, Canada	Swedish Board for Accreditation and Conformity Assessment (SWEDAC) (for mercury)
Slovenia	Seronorm whole blood level 1, NIST 1547, NIST 966, IAEA-086		
Czech Republic	Seronorm whole blood level 1		Czech Accreditation Institute (for trace elements by AAS in biological material)
Poland	Seronorm whole blood level 1	CDC, USA; Istituto Superiore de Sanita, Italy	
China	Seronorm whole blood level 1, 3, Dorm-2, Dolt-3, serum (ClinCheck)		

- non-exposed persons,
- people occupationally exposed to elemental mercury,
- “fish eaters.”

In a fourth intercomparison 3 different samples of fresh blood from the general population was used.

Figure 1 shows the results for these 6 samples when analyzed by the 5 laboratories that generated the data. If $|Z| \leq 2$ is applied, using a Z-score based on a target

relative standard deviation of 25% [18] as a criterion for acceptance, 4 out of the 5 laboratories reported satisfactory results for all samples, and 1 laboratory delivered 2 out of their 18 results outside the satisfactory range. The results of these intercomparisons thus showed results that could be considered as fair but the authors concluded that because samples from different countries were analyzed in different laboratories and systematic differences



On the horizontal axis: the mean result for all laboratory setups (up to 18, including also other laboratories than the here included) participating in the interlaboratory comparison.

On the vertical axis: the mean result for each of the 5 laboratories that carried out the analyses of blood in the present report. Six samples were analyzed, of which 3 were lyophilised (1 was a pooled sample from 12 non-exposed, 1 was from a worker with occupational exposure to inorganic mercury, and 1 was a pooled sample from 3 individuals with high fish consumption), and 3 samples were fresh blood from 3 healthy voluntary donors (data from Snoj Tratnik et al. [19]).

Figure 1. Results of interlaboratory comparisons between the 5 laboratories that carried out the analyses of blood in the present study on children aged 7–14 years, living in 2007–2008 in urban, rural, or potentially polluted areas in 6 European and 3 non-European countries: a) cadmium (Cd), b) lead (Pb), and c) mercury (Hg)

appeared between laboratories, these data could not be used for comparisons of exposure levels between countries [19]. Therefore, a subsample (urban children) from each country was analyzed centrally in 1 laboratory, as described in Hrubá et al. [9], where also comparisons between countries are reported. However, systematic differences between laboratories do not affect comparisons within each country. As a consequence, data are not invalidated for studies of regional variation within countries. The present report, which is based on the results from the 5 laboratories, is therefore devoted to comparisons within countries, not between countries.

Statistical analyses

Herein, only regional comparisons of exposure levels within countries were performed, thus, no comparisons were made between countries. The statistical analyses, as a consequence, concern regional differences within countries and exposure correlates other than geography. Assessment of regional differences between urban, rural and industrial areas were based on the ratios between rural and urban, and industrial and urban concentrations. These ratios were calculated based on data from the laboratory that had analyzed all samples from the specific country. First, due to skewed distributions, concentrations were transformed using the natural logarithm. Second, mean differences of log-transformed metal concentrations, corresponding to ratios in data not logarithmically transformed, were calculated and, for each country, estimates of the multiplicative effect of area (that is, the ratios rural/urban and industrial/urban concentrations) were presented. Statistical differences of the differences were evaluated using t-tests.

In order to find out if any geographical differences within the European countries could be ascribed to individual lifestyle factors, rather than differences in the environment, candidate correlates of Cd, Pb and Hg concentrations were evaluated for these countries. The analysis

included only the European countries, as these were expected to be fairly homogenous in models of exposure correlates, i.e., interaction between country and correlates should not occur for many of the correlates. The analysis was performed by first evaluating whether the relative difference between areas differed between countries, using log-linear models. As significant interactions between country and area were found for all 3 metals, country- and area-specific means were estimated. Thereafter, the potential association was analyzed between metal concentrations and the following 12 categorical variables, for which specific hypotheses had been formulated [9]: child's sex and age, parental education, smoking at home, type of water source, type of heating source, traffic density, the child's attempt to smoke, number of amalgam fillings, and intake of fish, shellfish, and offal. For those variables that were statistically significant, interactions between the significant influential variable and country, as well as interactions with living area, were also examined. Since a significant interaction between amalgam fillings and country was found, this relationship was studied separately for each country. Otherwise, the final model included country and area, and all other statistically significant correlates. "Statistically significant" denotes $p < 0.05$ (2-sided). Residual distributions were assessed for normality using q-q plots, and homogeneity of variance was assessed visually and using Levene's test. Influential data points were identified using the leverage measure. Stata 10 was used for all statistical analyses.

RESULTS

Cadmium

The variation within countries was substantial for B-Cd but there was no consistent pattern for rural areas, compared to urban ones (Table 3). A major impact of region was observed in Poland, where B-Cd in children for the selected industrial area was approx. double (2.1 times, 95% CI: 1.6–2.8) compared to urban children (geomet-

Table 3. Geometric means of children's blood metal concentrations (lead [Pb], mercury [Hg]) in urban areas of 9 countries and the relative difference comparing urban with rural and industrial areas

Country/Area/Pollution	Blood metal concentration					
	cadmium (Cd)		lead (Pb)		mercury (Hg)	
	central/local lab. [µg/l] (GM)	relative difference ^a point estimate (95% CI)	central/local lab. [µg/l] (GM)	relative difference ^a point estimate (95% CI)	central/local lab. [µg/l] (GM)	relative difference ^a point estimate (95% CI)
Croatia						
urban	0.17/0.26	1.00	17.9/19.4	1.00	n.a./0.44	1.00
rural		1.17 (1.07–1.27)*		1.13 (0.96–1.33)		0.82 (0.66–1.01)
industrial (petrochemical)		1.13 (1.03–1.23)*		1.18 (0.99–1.40)		0.65 (0.51–0.81)*
Czech Republic						
urban	0.13/0.18	1.00	15.5/24.3	1.00	0.21/0.42	1.00
rural		1.20 (1.03–1.42)*		0.62 (0.51–0.77)*		1.01 (0.75–1.37)
industrial (Pb)		1.60 (1.37–1.86)*		0.95 (0.78–1.16)		0.96 (0.72–1.28)
Poland						
urban	0.15/0.20	1.00	16.3/14.2	1.00	0.12/0.31	1.00
rural		1.20 (0.91–1.60)*		1.40 (1.18–1.67)*		0.70 (0.48–0.99)*
industrial (Pb)		2.13 (1.61–2.81)*		2.28 (1.92–2.70)*		0.57 (0.40–0.82)*
Slovakia						
urban	0.14/n.a.	1.00	19.4/n.a.	1.00	0.52/n.a.	1.00
rural		0.87 (0.77–0.99)*		1.12 (0.99–1.26)		0.52 (0.40–0.68)*
industrial (Pb)		0.99 (0.87–1.12)		1.82 (1.61–2.05)*		0.57 (0.44–0.74)*
Slovenia						
urban	0.14/0.27	1.00	13.4/20.2	1.00	n.a./0.94	1.00
rural		1.23 (1.09–1.38)*		1.10 (0.96–1.26)		0.76 (0.63–0.92)*
Industrial (Hg)		0.92 (0.82–1.04)		0.93 (0.81–1.07)		0.97 (0.80–1.17)
Sweden						
urban	0.11/n.a.	1.00	14.0/n.a.	1.00	0.43/n.a.	1.00
rural		0.84 (0.74–0.95)*		0.81 (0.69–0.95)*		1.40 (1.10–1.80)*
industrial (Pb)		1.01 (0.89–1.13)		1.03 (0.88–1.21)		2.09 (1.65–2.65)*
China						
urban	n.a./0.75	1.00	n.a./64.2	1.00	2.45/2.23	1.00
rural		0.95 (0.90–1.01)		0.72 (0.62–0.84)*		1.19 (1.07–1.34)*
industrial (Hg)		0.99 (0.93–1.05)		0.76 (0.64–0.90)*		3.07 (2.71–3.47)*
Ecuador						
urban	0.26/n.a.	1.00	31.7/n.a.	1.00	3.23/n.a.	1.00
rural		1.21 (1.05–1.40)*		1.26 (1.08–1.46)*		0.91 (0.74–1.12)
industrial (Hg)		1.04 (0.88–1.22)		0.95 (0.80–1.12)		1.50 (1.18–1.89)*

Table 3. Geometric means of children's blood metal concentrations (lead [Pb], mercury [Hg]) in urban areas of 9 countries and the relative difference comparing urban with rural and industrial areas – cont.

Country/Area/Pollution	Blood metal concentration					
	cadmium (Cd)		lead (Pb)		mercury (Hg)	
	central/local lab. [µg/l] (GM)	relative difference ^a point estimate (95% CI)	central/local lab. [µg/l] (GM)	relative difference ^a point estimate (95% CI)	central/local lab. [µg/l] (GM)	relative difference ^a point estimate (95% CI)
Morocco						
urban (Pb)	0.21/n.a.	1.00	71.0/n.a.	1.00	0.31/n.a.	1.00
rural		0.98 (0.83–1.15)		0.45 (0.37–0.54)*		0.52 (0.36–0.75)*
industrial (mix)		1.08 (0.88–1.33)		0.66 (0.51–0.84)*		1.25 (0.78–2.02)

GM – geometric mean; n.a. – not analyzed.

^a Between areas.

* The relative difference vs. the urban area is statistically significant ($p < 0.05$).

ric mean (GM) in central analyses of urban children in Poland: 0.15 µg/l, and in the Czech Republic 60% higher (1.6 times, 95% CI: 1.4–1.9, urban GM: 0.13 µg/l).

The only statistically significant correlation for B-Cd, except region, was traffic density (Table 4). Children who lived close to roads with a traffic density of ≥ 1 car/min had, on average, 7% higher levels than the others. The effect was consistent over the different European countries, i.e., no interaction with country was observed ($p = 0.2$). The effect did not remain statistically significant when urban children were excluded ($p = 0.15$). Traffic density, when included in the multivariate model, did not change the patterns of differences between urban/rural/industrial areas.

The B-Cd results obtained in the 3 local laboratories were all higher than in the central one, sometimes by a factor of 2 (Table 3).

Lead

A strong impact on the children's B-Pb from environmental contamination was seen in the selected industrial areas in Poland (2.3 times, 95% CI: 1.9–2.7, higher than in urban children, GM 16.3 µg/l) and Slovakia (1.8 times, 95% CI: 1.5–2.1, GM 19.4 µg/l) (Table 3). Within countries, B-Pb was higher in urban than in rural areas in

Sweden, the Czech Republic, China, and Morocco, though on quite different levels. On the other hand, rural children had statistically significantly higher B-Pb in both Poland and Ecuador, although the relative differences were small.

Boys had on average higher B-Pb than girls (Table 4). Also, exposure to environmental tobacco smoke correlated with B-Pb; children with such exposure had, on average, 10% higher concentrations. A similar relative increase was seen in children with frequent intake of offal. These factors were all consistent over all the European countries. When they were included in the multivariate model, they did not change the patterns of differences between areas.

For B-Pb, most – though not all – of the local laboratories gave higher results than the central one (Table 3).

Mercury

Children in rural areas could have either lower (Poland, Slovakia, Slovenia and Morocco) or higher (Sweden and China) B-Hg than children in the cities. Likewise, the children from industrial areas could have either lower (Croatia, Slovakia and Poland) or higher (Sweden, China, and Ecuador) B-Hg than the urban children (Table 3).

Table 4. Associations between metal concentrations in blood in children and potential determinants (data from the 6 European countries: Croatia, the Czech Republic, Poland, Slovakia, Slovenia, and Sweden)

Variable	Starting model ^a R ² [%]	Model with all influential variables included					
		relative change			interaction		
		point estimate	95% CI	p for trend	R ² [%]	with country p	with area p
Cadmium	57				60		
traffic density				0.014		0.2	0.4
<1 car/min		1.00	ref.				
≥1 car/min		1.07	1.01–1.12				
Lead	33				37		
sex				<0.001		0.8	0.6
girls		1.00	ref.				
boys		1.09	1.04–1.15				
smoking home				0.004		0.4	0.9
no smoker home		1.00	ref.				
a smoker home		1.10	1.03–1.18				
offal intake				0.004		0.9	0.2
≤1 meal/month		1.00	ref.				
>1 meal/month		1.09	1.03–1.16				
Mercury	36				40		
fish intake				<0.001		0.5	0.7
<1 meal/month		1.00	ref.				
1–3 meals/month		1.10	0.99–1.22				
>3 meals/month		1.39	1.24–1.55				
amalgam fillings (Table 5)				<0.001		0.026	0.7

R² – explained variance, expressed as percentage of the total variance.

^a Starting model includes country, area, and interaction term between country and area.

Bold is a statistically significant value (p < 0.05).

As expected, B-Hg increased with rising intake of fish (Table 4, consistent for the different European countries) and number of amalgam fillings (Table 5, varying effect in different countries). However, these factors, when included in the multivariate model, did not change the patterns of differences between areas.

Two of the 3 countries that had B-Hg determined both locally and in a central laboratory showed considerably higher concentrations than the central laboratory (Table 3).

DISCUSSION

Regional variation

The results show that children in certain contaminated areas, so called “hot spots,” in Europe are considerably more exposed to cadmium and/or lead than children living in other areas. The possibility that the exposure at these “hot spots” has decreased since the sampling in 2007–2008 can not be excluded but there is no indications of that.

Table 5. Associations between mercury concentrations in blood and amalgam fillings, and interaction with country (data from 6 European countries) in the study on children aged 7–14 years, living in 2007–2008 in urban, rural, or potentially polluted areas

Amalgam fillings	Mercury concentrations association				
	model with area R ² [%]	model with area and fish intake ^a			R ² [%]
		relative change		p	
		point estimate	95% CI		
0–1		1.00	ref.		
≥2					
Croatia	7	1.27	1.04–1.56	0.019	14
Czech Republic	0	1.15	0.92–1.45	0.23	5
Poland	6	1.66	0.93–2.96	0.090	8
Slovakia	15	1.78	1.43–2.20	<0.001	32
Slovenia	5	1.19	1.02–1.42	0.038	8
Sweden	20	— ^b			

R² – explained variance, expressed as percentage of the total variance.

^a Fish intake was an influential variable (Table 4).

^b No child with ≥2 amalgam fillings.

Bold are statistically significant values (p < 0.05).

Contamination of the Eastern European “hot spots” largely stems from old heavy metallurgic industry that expanded largely during the 1950s and 1960s. Examples of children’s exposure from such areas have been published before, for example from Poland, where living in the vicinity of a metallurgical slag heap added significantly to the children’s lead exposure [13]. However, this industrial expansion does not seem to have produced significant nation-wide exposures, as B-Pb and B-Cd in urban children varied only little between the European countries studied herein [9]. Taken together, this means that exposures due to local contamination is a significant problem in certain areas in Europe. The extent of this problem is unknown.

In China and Ecuador, B-Hg was higher than in the European countries. The industrially contaminated area studied in China is a rather special one, with Hg mines. Evidently, the Hg from this source exposed not only workers, but also children living in the area. In Ecuador, a similar situation was evident for children from the gold-mining

area, where large amounts of Hg were used to refine gold, sometimes even indoors at home close to children.

Some relatively small regional differences were noted, for example the somewhat higher B-Pb in rural areas of Ecuador and Poland vs. urban areas. The causes are unknown to the authors. Possibly, differences in pollution control of drinking water (e.g. from private wells), in lifestyle not captured by the questionnaire, or proximity to industrial sources could play roles.

Other correlates of exposure

Blood cadmium was associated with traffic density, confirming previous finding [9]. The cause is obscure, but may be due to Cd in tyres, bioavailability of Cd in freshly generated combustion particles, or residual confounding from socioeconomic factors or different dietary habits. The B-Cd was not associated with other factors such as consumption of offal, fish or shellfish. Consumption of other dietary items such as rice, known to influence B-Cd, was not considered, but it appears unlikely that

traffic density was confounded by rice consumption. Therefore, the association between B-Cd and traffic density may well be causal and not caused by confounding. Neither second-hand smoking, nor age was associated with B-Cd. For second-hand smoke, this is in line with other data [20,21].

As to B-Pb, leaded petrol was still relevant in countries with recent (China) or probably prevailing (Morocco) use, where the urban children showed higher levels than the children from other regions. This is suggestive of petrol-Pb exposure or, possibly, Pb in the local water pipes [5]. Alternative explanations in the Chinese city could be combustion of coal [5,22] or industrial emissions [23].

In accordance with a wealth of previous findings, male sex [24] and exposure to environmental tobacco smoke [20] are established determinants for B-Pb. Moreover, there was an association with offal intake. Hence, consumption of, e.g., liver and kidney may contribute to the Pb intake in children, in the same way as has been shown in adults [25]. However, a previously reported by the authors relation to shellfish [9], was not confirmed.

For B-Hg, the relations to fish intake and amalgam fillings were expected, on the basis of earlier observations in adults [26,27] and children [9]. However, it is remarkable that even the rather small intake of fish in these children, as well as their very low number of amalgam fillings, were influential. As regards mercury exposure in Europe, it should be noted that populations in several European countries with coastal regions and larger fish consumption show higher mercury levels than in the countries here represented [28].

Taken together, the observed correlates, other than region, made only a minor contribution (an additional 3–4%) to the explained variance in the statistical model, which probably means that their impact on exposure is small in relation to the dominating factors, i.e., geography, which accounted for 33–57% of the variance.

Design issues

A limitation is that a relatively low number of children from each area was examined. Another limitation is that sampling was carried out only in single schools, and the results cannot give a full picture of the country, or even the specific area. An asset of the present study is the use of common sampling procedures and questionnaires/interviews, but a limitation is that quite simple questionnaires were used. This means that identification of correlates (potential determinants) can only be made in a relatively rough way. Further, soil, air, food or water concentrations of the 3 metals were not investigated.

Blood was used for the analyses. Blood requires invasive sampling but is for lead the recommended matrix. It is also a useful matrix for biomonitoring of cadmium and mercury exposure, although urine is often chosen for assessment of cadmium exposure, especially long-term (several years) exposure in adults. In contrast, B-Cd reflects a combination of long- and short-term exposure. Further, urinary biomonitoring of cadmium involves difficulties as regards adjustment for dilution, urinary flow, and co-elution with other substances, for example proteins [29]. As regards children, very little is known about these effects. Therefore, B-Cd is considered as a good alternative for a comparison of cadmium exposure among children from different regions. A limitation related to the above is that for cadmium it is difficult to compare the present data with studies based on urine samples. It should also be noted that sampling was made in 2007–2008. The authors do believe that the observations of higher exposures in certain polluted areas in Europe are still relevant, as there are no strong indications of general improvements lately as regards “hot spots” exposures.

A somewhat complicated limitation is that the authors did not have full comparability between countries, because different laboratories were used for the analyses of samples from the various countries. It is then hard to make conclusions on differences between countries. Any observations will raise the question: do results reflect variation in actual

concentrations or just in laboratory performance? Given the modest variation between countries, an extremely good agreement between laboratories would be required; analyses of the same samples should have to fall within roughly $\pm 20\%$ or even less, considering that variation within $\pm 20\%$ means that the highest acceptable value ($1.2 \times$ target concentration) will be 1.5 times the lowest acceptable value ($0.8 \times$ target concentration). This quality control study did not show such low variation. That is the reason why some of the samples were re-analyzed in a single laboratory before an international comparison was made [9]. By the same reason, the authors here refrain from any international comparisons of concentrations and choose not to tabulate each regions concentration in Table 3, as such tabulation might lead readers to make inconclusive comparisons. The table instead shows ratios vs. the urban results. In this way, data from different laboratories can be used for combined analyses. Thus, while direct international comparisons are hindered, conclusions can still be made about regional variation and correlations with exposure other than geography.

It should be noted that difficulties with comparability between laboratories is a common limitation that is hard to overcome. For example, the quality control program applied in the European study of cadmium in urine [12], after taking into account what was technically feasible and realistic in current routine practice, allowed as much as $\pm 50\%$ variation in its interlaboratory comparison ($|Z| \leq 2$, using a Z-score based on a target relative standard deviation of 25%; [18]). This problem of comparability between laboratories is a likely explanation of the fact that there are remarkably few international coherent studies that have presented data on toxic metals that are fully comparable between countries. As regards studies of both B-Pb and B-Cd the authors know of only 3 such studies that include >2 countries [9,10,30] despite significant biomonitoring efforts in, for example, Europe [31].

All European countries' data were modelled together. Then there is a risk that differences in laboratory performance

may invalidate analyses, (e.g., factors correlating with blood concentrations) (Table 4). The authors therefore included country in the model, which accounts for most calibration errors. In this way the combined dataset could be used for analysis of factors correlating with exposure.

Health aspects

The B-Cd in the presently studied children is generally low, but because of the long-term accumulation of Cd in the body, this exposure may still lead to risks of kidney and bone effects later in life, in particular in females in contaminated areas [1,7,32]. Possibly, these exposure levels can also affect cognitive development [6].

The B-Pb in the industrial areas in both Poland and Slovakia, and in all Ecuadorian regions, showed geometric means of about $35 \mu\text{g/l}$, with a considerable fraction above $50 \mu\text{g/l}$, a level nowadays regarded as a cause of an increased risk of slight impairments of cognitive functions [2,3,5]. The levels in China and Morocco were even higher. Recent risk assessments have not ascertained a safe level. In the USA, a "reference level" of $50 \mu\text{g/l}$ is used to "identify children who have been exposed to lead and who require case management" ([33]; recently suggested to be lowered to $35 \mu\text{g/l}$ [34]) and European Food Safety Authority [2] and World Health Organization and Food and Agriculture Organization [3] jointly proposed a tolerable intake of lead corresponding to a B-Pb of $12 \mu\text{g/l}$.

The B-Hg in the European and Moroccan children was lower than those known for long to cause central nervous and kidney effects in adults [4,26,27]. However, higher levels occur in the areas studied in China and Ecuador. The highest observed B-Hg may cause a risk of damage to the fetal central nervous system, if retained into the fertile period in women, especially if it is in the form of methyl-Hg. High exposure to methyl-Hg is a possibility [35,36] but the present study could not differentiate between inorganic Hg, from e.g. inhalation of elemental mercury, and methyl-Hg from contaminated food.

Recommendations

With the present blood metal concentrations and the knowledge of their toxic effects, there is a need for prevention programs in Europe that decrease human exposure in industrially Cd- and Pb-contaminated areas. Then, the strategy should be to specifically act on “hot spots” exposures, which are unsafe, identifiable, and preventable. Also in the countries outside Europe, measures need to be taken to reduce exposure to all 3 metals among children. Prevention programs should result in clean-up of areas, and/or alteration of industrial activity. Human biomonitoring of exposure is then a very useful component in the identification and prioritization of areas.

CONCLUSIONS

These mainly European results indicate that some children experience about doubled exposures to toxic elements just because of where they live. These exposures are unsafe, identifiable, and preventable and therefore call for preventive actions.

In addition to previously well-known determinants, traffic density appears to be associated with children's B-Cd, though with a relatively modest effect.

ACKNOWLEDGMENTS

The authors acknowledge contributions of late Mr. Mladen Kršnik and late Dr. Raúl Harari to this study. The authors thank Prof. Ulf Strömberg, University of Gothenburg, for advice on statistical methods, and Dr. Nils-Göran Lundström, Umeå University, for project administrative support. Technical assistance was given by Ms Anna Akantis and Mr Giovanni Ferrari. The authors also acknowledge all the participating children, and those who assisted in the field work.

Author contributions

Research concept: Staffan Skerfving, Ingvar A. Bergdahl

Research methodology: Františka Hrubá, Milena Černá, Chunying Chen, Florencia Harari, Milena Horvat,

Kvetoslava Koppová, Andrea Krsková, Jawhar Laamech, Yu-Feng Li, Lina Löfmark, Thomas Lundh, Badiia Lyoussi, Darja Mazej, Joško Osredkar, Krystyna Pawlas, Natalia Pawlas, Adam Prokopowicz, Gerda Rentschler, Janja Snoj Tratnik, Johan Sommar, Věra Spěváčková, Zdravko Špirić, Staffan Skerfving, Ingvar A. Bergdahl

Collecting material: Milena Černá, Chunying Chen, Florencia Harari, Milena Horvat, Kvetoslava Koppová, Andrea Krsková, Jawhar Laamech, Yu-Feng Li, Lina Löfmark, Thomas Lundh, Badiia Lyoussi, Darja Mazej, Joško Osredkar, Krystyna Pawlas, Natalia Pawlas, Adam Prokopowicz, Gerda Rentschler, Janja Snoj Tratnik, Věra Spěváčková, Zdravko Špirić, Staffan Skerfving

Statistical analysis: Františka Hrubá, Johan Sommar, Staffan Skerfving, Ingvar A. Bergdahl

Interpretation of results: Františka Hrubá, Milena Černá, Chunying Chen, Florencia Harari, Milena Horvat, Kvetoslava Koppová, Andrea Krsková, Jawhar Laamech, Yu-Feng Li, Lina Löfmark, Thomas Lundh, Badiia Lyoussi, Darja Mazej, Joško Osredkar, Krystyna Pawlas, Natalia Pawlas, Adam Prokopowicz, Gerda Rentschler, Janja Snoj Tratnik, Johan Sommar, Věra Spěváčková, Zdravko Špirić, Staffan Skerfving, Ingvar A. Bergdahl

REFERENCES

1. EFSA. European Food Safety Authority. Cadmium in food - scientific opinion of the Panel on Contaminants in the food chain. *EFSA J.* 2009;7(3):980. <https://doi.org/10.2903/j.efsa.2009.980>.
2. EFSA. European Food Safety Authority. Scientific opinion on lead in food. *EFSA J.* 2010;8(4):1570. <https://doi.org/10.2903/j.efsa.2010.1570>.
3. JECFA. Joint FAO/WHO Expert Committee on Food Additives and Contaminants. Evaluation of certain food additives and contaminants. Seventy-third report of the Joint FAO/WHO Expert Committee on Food Additives and Contaminants. Technical Report Series No. 960. Geneva, World Health Organization; 2011.

4. Berlin M, Zalups RK, Fowler BA. Mercury. In: Nordberg GF, Fowler BA, Nordberg M, editors. Handbook on the toxicology of metals, Volume II. Fourth edition. Academic Press ISBN 978-0-12-398293-3; 2014. p. 1013-75.
5. Bergdahl IA, Skerfving S. Lead. In: Nordberg GF, Costa M (editors) Handbook on the toxicology of metals, Volume II, Fifth edition. Academic Press. ISBN 978-0-12-922946-0; 2022. p. 427-93. <https://doi.org/10.1016/B978-0-12-822946-0.00036-2>.
6. Gustin K, Tofail F, Vahter M, Kippler M. Cadmium exposure and cognitive abilities and behavior at 10 years of age: A prospective cohort study. *Environ Int.* 2018;113:259-68. <https://doi.org/10.1016/j.envint.2018.02.020>.
7. Nordberg GF, Bernard A, Diamond GL, Duffus JH, Illing P, Nordberg M, et al. Risk assessment of effects of cadmium on human health (IUPAC Technical Report). *Pure Appl Chem.* 2018;90(4):755-808. <https://doi.org/10.1515/pac-2016-0910>.
8. EFSA. European Food Safety Authority. Scientific opinion on the risk for public health to the presence of mercury and methylmercury in food. *EFSA J.* 2012;10(12):2985. <https://doi.org/10.2903/j.efsa.2012.2985>.
9. Hrubá F, Strömberg U, Černá M, Chen C, Harari F, Harari R, et al. Blood cadmium, mercury, and lead in children: an international comparison of cities in six European countries, and China, Ecuador, and Morocco. *Environ Int.* 2012;41:29-34. <https://doi.org/10.1016/j.envint.2011.12.001>.
10. Pawlas N, Strömberg U, Carlberg B, Cerna M, Harari F, Harari R, et al. Cadmium, mercury and lead in the blood of urban women in Croatia, the Czech Republic, Poland, Slovakia, Slovenia, Sweden, China, Ecuador and Morocco. *Int J Occup Med Environ Health.* 2013;26(1):58-72. <https://doi.org/10.2478/S13382-013-0071-9>.
11. Rentschler G, Rodushkin I, Cerna M, Chen C, Harari F, Harari R, et al. Platinum, palladium, rhodium, molybdenum and strontium in blood of urban women in nine countries. *Int J Hyg Environ Health.* 2018;221(2):223-30. <https://doi.org/10.1016/j.ijheh.2017.10.017>.
12. Snoj Tratnik J, Kocman D, Horvat M, Andersson AM, Juul A, Jacobsen E, et al. Cadmium exposure in adults across Europe: Results from the HBM4EU Aligned Studies survey 2014-2020. *Int J Hyg Environ Health.* 2022 Sep;246:114050. <https://doi.org/10.1016/j.ijheh.2022.114050>.
13. Kowalska M, Kulka E, Jarosz W, Kowalski M. The determinants of lead and cadmium blood levels for preschool children from industrially contaminated sites in Poland. *Int J Occup Med Environ Health.* 2018;31:351-359. <https://doi.org/10.13075/ijomeh.1896.01153>.
14. Laamech J, Bernard A, Dumont X, Benazzouz B, Lyoussi B. Blood lead, cadmium and mercury among children from urban, industrial and rural areas of Fez Boulemane region (Morocco): relevant factors and early renal effects. *Int J Occup Med Environ Health.* 2014;27(4):641-59. <https://doi.org/10.2478/s13382-014-0275-7>.
15. Barany E, Bergdahl IA, Schütz A, Skerfving S, Oskarsson A. Inductively coupled plasma mass spectrometry for direct multi-element analysis of diluted human blood and serum. *J Anal At Spectrom.* 1997;12(9):1005-9. <https://doi.org/10.1039/a700904f>.
16. Propkowicz A, Pawlas N, Ochota P, Szuła M, Sobczak A, Pawlas K. Blood Levels of Lead, Cadmium, and Mercury in Healthy Women in their 50s in an Urban Area of Poland: A Pilot Study. *Pol J Environ Stud.* 2014;23(1):167-75.
17. Sandborgh-Englund G, Elinder CG, Langworth S, Schütz A, Ekstrand J. Mercury in biological fluids after amalgam removal. *J Dent Res.* 1998;77(4):615-24. <https://doi.org/10.1177/00220345980770041501>.
18. Nübler S, López ME, Castaño A, Mol H, Schäfer M, Haji-Abbas-Zarrabi K, et al. Interlaboratory comparison investigations (ICI) and external quality assurance schemes (EQUAS) for cadmium in urine and blood: Results from the HBM4EU project. *Int J Hyg Environ Health.* 2021;234:113711. <https://doi.org/10.1016/j.ijheh.2021.113711>.
19. Snoj Tratnik J, Mazej D, Horvat M. Analytical quality requirements in human biomonitoring programs: Trace elements in human blood. *Int J Environ Res Public Health.* 2019;16(13):2287. <https://doi.org/10.3390/ijerph16132287>.

20. Willers S, Schütz A, Attewell R, Skerfving S. Relation between lead and cadmium in blood and the involuntary smoking of children. *Scand J Work Environ Health*. 1989; 14(6):385-9. <https://doi.org/10.5271/sjweh.1905>.
21. Richter PA, Bishop EE, Wang J, Swahn MH. Tobacco smoke exposure and levels of urinary metals in the U.S. youth and adult population: the National Health and Nutrition Examination Survey (NHANES) 1999-2004. *Int J Environ Res Public Health*. 2009;6(7):1930-46. <https://doi.org/10.3390/ijerph6071930>.
22. Liang F, Zhang G, Tan M, Yan C, Li X, Li Y, et al. Lead in children's blood is mainly caused by coal-fired ash after phasing out of leaded gasoline in Shanghai. *Environ Sci Technol*. 2010;44(12):4760-5. <https://doi.org/10.1021/es9039665>.
23. Zhao ZQ, Zhang W, Li XD, Yang Z, Zheng HY, Ding H, et al. Atmospheric lead in urban Guiyang, Southwest China: Isotope source signatures. *Atmos Environ*. 2015;115:163-9. <https://doi.org/10.1016/j.atmosenv.2015.05.049>.
24. Skerfving S, Bencko V, Vahter M, Schütz A, Gerhardsson L. Environmental health in the Baltic region – toxic metals. *Scand J Work Environ Health*. 1999;25 Suppl 3:40-64.
25. Bensryd I, Rylander L, Högstedt B, Aprea P, Bratt I, Fåhræus C, et al. Effect of acid precipitation on retention and excretion of elements in man. *Sci Total Environ*. 1994;145(1-2):81-102. [https://doi.org/10.1016/0048-9697\(94\)90299-2](https://doi.org/10.1016/0048-9697(94)90299-2).
26. WHO. Environmental Health Criteria 101. Methylmercury. Geneva: International Programme on Chemical Safety, World Health Organization; 1990.
27. WHO. Environmental Health Criteria 118. Inorganic mercury. Geneva: International Programme on Chemical Safety, World Health Organization; 1991.
28. Visnjevec AM, Kocman D, Horvat M. Human mercury exposure and effects in Europe. *Environ Toxicol Chem* 2014;33(6):1259-70. <https://doi.org/10.1002/etc.2482>.
29. Akerstrom M, Barregard L, Lundh T, Sallsten G. Variability of urinary cadmium excretion in spot urine samples, first morning voids, and 24 h urine in a healthy non-smoking population: implications for study design. *J Expo Sci Environ Epidemiol*. 2014;24(2):171-9. <https://doi.org/10.1038/jes.2013.58>.
30. Vahter M, Slorach S. Exposure monitoring of lead and cadmium. An international pilot study within the UNEP/WHO human exposure assessment location (HEAL) project. Technical report, WHO, Nairobi; 1991.
31. Joas A, Knudsen LE, Kolossa-Gehring M, Sepai O, Castleyn L, Schoeters G, et al. Policy recommendations and cost implications for a more sustainable framework for European human biomonitoring surveys. *Environ Res*. 2015;141:42-57. <https://doi.org/10.1016/j.envres.2014.10.012>.
32. Åkesson A, Barregard L, Bergdahl IA, Nordberg GF, Nordberg M, Skerfving S. Non-renal effects and the risk assessment of environmental cadmium exposure. *Environ Health Perspect*. 2014;122(5):431-8. <https://doi.org/10.1289/ehp.1307110>.
33. U.S. CDC [Internet]. CDC Response to Advisory Committee on Childhood Lead Poisoning Prevention. Recommendations in “Low Level Lead Exposure Harms Children: A Renewed Call of Primary Prevention”. 2021. Available from: https://www.cdc.gov/nceh/lead/acclpp/cdc_response_lead_exposure_recs.pdf. Accessed 4 September 2021
34. LEPAC [Internet]. Centers for Disease Control and Prevention Lead Exposure and Prevention Advisory Committee (LEPAC). Meeting held via Zoom web video conferencing May 14, 2021 9:00 A.M. Available from: <https://www.cdc.gov/nceh/lead/advisory/lepac-meeting-5-14-21.html>. Accessed 4 September 2021
35. Li P, Li Y, Feng X. Mercury and selenium interactions in human blood in the Wanshan mercury mining area, China. *Sci Tot Environ*. 2016;573:376-81. <https://doi.org/10.1016/j.scitotenv.2016.08.098>.
36. Li P, Guo S, Zhao J, Gao Y, Li YF. Human biological monitoring of mercury through hair samples in China. *Bull Environ Contam Toxicol*. 2019;102(5):701-7. <https://doi.org/10.1007/s00128-019-02563-8>.