

Errata for thesis:

Martin Angelin (2015) **Travel – a risk factor for disease and spread of antibiotic resistance**. Department of Clinical Microbiology, Infectious Diseases, Umeå University.

ISBN 978-91-7601-348-9

ISSN 0346-6612

New Series No: 1754

Corrections

Page 7, Table 1:

Polio	Case reports Last case of imported Polio in industrialized countries was in 2007. Recent spread from Iraq and Syria to East Africa.	High
Tuberculosis	1.28-2.8/1000 person-months of travel with PPD conversion, <u>0.06-0.6/1000</u> person-months of travel with active infection	Low in adults

Page 15, lines 13-15: Integrons are an important part of HGT and since they cannot be transmitted independently they are ~~although they may be transmitted independently, they are most often~~ found on plasmids as well as form parts of transposons.

Page 16, lines 35-36: (...), such as *Escherichia coli*, *Klebsiella pneumoniae* and *Proteus mirabilis*.

Page 20, lines 10-11: Several risk factors for infection with ESBL-PE producing bacteria have been identified,(...)

Page 29, lines 5-8: We aimed for a very high sequence coverage (number of times each base pair is covered when aligning sequence reads) to also find less abundant resistance genes.

Page 32, lines 26-27: As with hygiene advice, most travel-related advice is experience based rather than research based, but attaining true research-based advice is difficult.

Page 34, lines 15-16: Experience-based advice does exist and needs to be utilised by travellers.